Phospho-eIF4E (Ser209) Ab

Cat.#: AF3110 Concn.: 1mg/ml Mol.Wt.: 25kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-elF4E (Ser209) Ab detects endogenous levels of

eIF4E only when phosphorylated at Serine 209

Immunogen: A synthesized peptide derived from human eIF4E around the

phosphorylation site of Serine 209

Uniprot: P06730

Description: eIF4E a protein of the eukaryotic initiation factor 4E family.

Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome binding by inducing the

unwinding of the mRNAs secondary structures.

Similarity: Belongs to the eukaryotic initiation factor 4E family.

Storage Condition and

Buffer:


Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt



Western blot analysis of eIF4E phosphorylation expression in FBS treated NIH-3T3 whole cell lysates,The lane on the left is

treated with the antigen-specific peptide.




Western blot analysis of Phospho-eIF4E (Ser209) expression in various lysates



Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com



AF3110 at 1/200 staining human colon carcinoma tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat antirabbit Ab was used as the secondary.



AF3110 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.