


53BP1 (Phospho-Ser1778) Ab

Cat.#: AF8337 Size: 50ul,100ul,200ul	Concn.: 1mg/ml Source: Rabbit	Mol.Wt.: 214 kDa Clonality: Polyclonal
Application:	WB 1:1000-3000, IF/ICC 1:100-1:500	
Reactivity:	Human	
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.	
Specificity:	53BP1 (Phospho-Ser1778) Ab detects endogenous levels of 53BP1 only when phosphorylated at Ser1778	
Immunogen:	A synthesized peptide derived from human 53BP1 (Phospho-Ser1778)	
Uniprot:	Q12888	
Subcellular Location:	Nucleus. Chromosome > centro Associated with kinetochores. B in some cells. Recruited to sites double stand breaks. Methylatic required for efficient localization	oth nuclear and cytoplasmic of DNA damage, such as on of histone H4 at 'Lys-20' is
Similarity:	The Tudor-like region mediates dimethylated at 'Lys-20' (H4K20 Interaction with NUDT16L1/TIRF domain and prevents recruitme (PubMed:28241136).The UDR (u recruitment) motif specifically r H2A monoubiquitinated at 'Lys- (PubMed:23760478, PubMed:24 the UDR blocks interaction with (PubMed:24703952).	Ome2) (PubMed:17190600). R masks the Tudor-like nt to chromatin ubiquitin-dependent ecognizes and binds histone 15' (H2AK15ub) P03952). Phosphorylation of
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt	



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis 53BP1 (Phospho-Ser1778) using UV treated Jurkat whole cell lysates



AF8337 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.