


Cyclin E1 (Phospho-Ser399) Ab

Cat.#: AF8341
Size: 50ul,100ul,200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 49kDa
Clonality: Polyclonal

Application:	WB 1:1000-3000, IF/ICC 1:100-1:500
Reactivity:	Human
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Specificity:	Cyclin E1 (Phospho-Ser399) Ab detects endogenous levels of Cyclin E1 only when phosphorylated at Ser399
Immunogen:	A synthesized peptide derived from human Cyclin E1 (Phospho-Ser399)
Uniprot:	P24864
Subcellular Location:	Nucleus.
Tissue Specificity:	Highly expressed in testis and placenta. Low levels in bronchial epithelial cells.
Similarity:	Belongs to the cyclin family. Cyclin E subfamily.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt



Western blot analysis Cyclin E1 (Phospho-Ser399) using LPS treated HeLa whole cell lysates



AF8341 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.