APC970Hu01 100µg Active Cathepsin V (CTSV) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1th Edition (Apr, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Gly64~Val334

Tags: N-terminal His-tag

Purity: >95%

Buffer Formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.8

Predicted Molecular Mass: 31.1kDa

Accurate Molecular Mass: 32kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

GEYSQGK HGFTMAMNAF GDMTNEEFRQ MMGCFRNQKF RKGKVFREPL FLDLPKSVDW RKKGYVTPVK NQKQCGSCWA FSATGALEGQ MFRKTGKLVS LSEQNLVDCS RPQGNQGCNG GFMARAFQYV KENGGLDSEE SYPYVAVDEI CKYRPENSVA NDTGFTVVAP GKEKALMKAV ATVGPISVAM DAGHSSFQFY KSGIYFEPDC SSKNLDHGVL VVGYGFEGAN SNNSKYWLVK NSWGPEWGSN GYVKIAKDKN NHCGIATAAS YPNV [ACTIVITY]

Cathepsin V (CTSV) is a lysosomal cysteine proteinase which belongs to peptidase C1 family. It is expressed in normal human thymus, testis and corneal epithelium. Cathepsin V plays an important role in corneal physiology and mediates degradation of invariant chain in human thymus. Besides. Retinoblastoma Protein 1 (RB1) has been identified as an interactor of CTSV, thus a binding ELISA assay was conducted to detect the interaction of recombinant human CTSV and recombinant human RB1. Briefly, CTSV were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to RB1-coated microtiter wells and incubated for 2h at 37℃. Wells were washed with PBST and incubated for 1h with anti-CTSV pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of CTSV and RB1 was shown in Figure 1, and this effect was in a dose dependent manner.




Figure 1. The binding activity of CTSV with RB1.

[IDENTIFICATION]





Figure 2. SDS-PAGE

Sample: Active recombinant CTSV, Human





Sample: Recombinant CTSV, Human;

Antibody: Rabbit Anti-Human CTSV Ab (PAC970Hu01)