## **GAS6 Antibody**

Cat.#: DF8659 Concn.: 1mg/ml Mol.Wt.: 79 kDa,70kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500,

ELISA(peptide) 1:20000-1:40000

\*The optimal dilutions should be determined by the end

user.

Reactivity: Human, Mouse

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: GAS6 Antibody detects endogenous levels of total GAS6.

Immunogen: A synthesized peptide derived from human GAS6,

corresponding to a region within the internal amino acids.

Uniprot: Q14393

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from MCF7, using GAS6 Antibody. The lane on the left was treated with blocking peptide.

Observed bands: 70 kDa.


Western blot analysis GAS6 using RAW264.7 whole cell

Ivsates




## **Affinity Biosciences**

website:www.affbiotech.com order:order@affbiotech.com



DF8659 at 1/100 staining Human melanoma tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.



DF8659 staining Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF8659 1:200) and mouse antibeta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.