

MAA091Hu21 Monoclonal Antibody to Macrophage Derived Chemokine (MDC) Organism Species: Homo sapiens (Human) *Instruction manual* 

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Suite 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.us | mail@cloud-clone.as Export Processing Zone, Wuhan, Hubei 436056, PRC | 0086-000-880-0687 | www.cloud-clone.com | mail@cloud-clone.com

# Cloud-Clone Corp.

### [PROPERTIES]

Source: Monoclonal antibody preparation Host: Mouse Antibody isotype: IgG2b Kappa Purification: Protein A/G Affinity Chromatography. Clone number: D1 Traits: Liquid Concentration: 500µg/mL UOM: 10µg Applications: WB; IHC; ICC; IP.

#### [IMMUNOGEN]

Immunogen: Recombinant MDC (Gly25~Gln93) expressed in *E.coli*. Accession No.: RPA091Hu01

### [APPLICATIONS]

Western blotting: 0.5-5µg/mL Immunocytochemistry in formalin fixed cells: 5-30µg/mL Immunohistochemistry in formalin fixed frozen section: 5-30µg/mL Immunohistochemistry in paraffin section: 5-30µg/mL Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the

## Coud-Clone Corp.

protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [IDENTIFICATION]




Figure 1. Western Blot Sample: Recombinant MDC, Human