MAS085Mi21

Monoclonal Antibody to Angiotensin 1-7 (Ang1-7) Organism Species: Homo sapiens (Human) Mus musculus (Mouse) Rattus norvegicus (Rat) Equus caballus; Equine (Horse) Ovis aries; Ovine (Sheep) Canis familiaris; Canine (Dog) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Soite 226, Houston, TX 77094, USA | 001-888-960-7402 | www.cloud-cloue.ox | mail@cloud-clone.ax Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0687 | www.cloud-clone.com | mail@cloud-clone.com

Cloud-Clone Corp.

[PROPERTIES]

Source: Monoclonal antibody preparation Host: Mouse Antibody isotype: IgG2b Kappa Purification: Protein A/G Affinity Chromatography. Clone number: A3 Traits: Liquid Concentration: 500µg/mL UOM: 200µg Applications: ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

[IMMUNOGEN]

Immunogen: CPS085Mi11-BSA conjugated Angiotensin 1-7 (Ang1-7) Immunogen: Synthetic Peptide, Ang1-7 conjugated to OVA Accession No.: CPS085Mi21 Sequence: The target peptide sequence is listed below. DRVYIHP

[ORGANISM SPECIES MORE]

React with: human, mouse, rat, equine, canine, ovine; Other species have not been detected.

[APPLICATIONS]

Immunocytochemistry in formalin fixed cells: 5-30ug/ml Immunohistochemistry in formalin fixed frozen section: 5-30ug/ml Immunohistochemistry in paraffin section: 5-30ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.