

PAA143Rb51 Polyclonal Antibody to Vascular Endothelial Growth Factor A (VEGFA) Organism Species: Oryctolagus cuniculus (Rabbit) Instruction manual

> FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

> > 12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Soite 226, Houston, TX 77094, USA | 001-888-960-7402 | www.choud-choue.ox | mail@choud-chone.as Export Processing Zone, Wuhan, Hubei 436056, PRC | 0086-000-880-0687 | www.choud-clone.com | mail@choud-chone.com

Cloud-Clone Corp.

[PROPERTIES]

Source: Polyclonal antibody preparation Host: Cavia Purification: Antigen-specific Affinity Chromatography. Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

[IMMUNOGEN]

Immunogen: Recombinant VEGFA (Leu1~Lys120) expressed in *E.coli*. Accession No.: RPA143Rb01

[APPLICATIONS]

Western blotting: 0.5-2ug/ml Immunocytochemistry in formalin fixed cells: 5-20ug/ml Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant VEGFA disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Coud-Clone Corp.

Loading Buffer: 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN₃.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. Western Blot Sample: Recombinant VEGFA, Rabbit