

PAA357Hu01
Polyclonal Antibody to Fibroblast Growth Factor 5 (FGF5)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

# [ PRODUCT INFORMATION ]

Immunogen: FGF5, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant FGF5 (Glu23~Gly268) expressed in *E.coli*.

Accession No.: RPA357Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- EKRLAPKG QPGPAATDRN PRGSSSRQSS SSAMSSSSAS SSPAASLGSQ GSGLEQSSFQ WSPSGRRTGS LYCRVGIGFH LQIYPDGKVN GSHEANMLSV LEIFAVSQGI VGIRGVFSNK FLAMSKKGKL HASAKFTDDC KFRERFQENS YNTYASAIHR TEKTGREWYV ALNKRGKAKR GCSPRVKPQH ISTHFLPRFK QSEQPELSFT VTVPEKKKPP SPIKPKIPLS APRKNTNSVK YRLKFRFG

9th Edition (Revised in Jul, 2013)

Sample: Recombinant FGF5, Human



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against FGF5. It has been selected for its ability to recognize FGF5 in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant FGF5 (Glu23~Gly268) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.