## Coud-Clone Corp.

#### PAA797Mu01 Polyclonal Antibody to Angiotensinogen (AGT) Organism Species: Mus musculus (Mouse) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [PRODUCT INFORMATION]

Immunogen: AGT, Mouse Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant AGT (Ala23~Val477) expressed in E.coli.

USCN Accession No.: RPA797Mu01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and S-tag and its sequence is listed below.

MHHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF-AGDRVYIH PFHLLYHNKS TCAQLENPSV ETLPESTFEP VPIQAKTSPV NEKTLHDQLV LAAEKLEDED RKRAAQVAMI ANFVGFRMYK MLNEAGSGAS GAILSPPALF GTLVSFYLGS LDPTASQLQT LLDVPVKEGD CTSRLDGHKV LAALRAVQGL LVTQGGSSSQ TPLLQSIMVG LFTAPGFRLK HSFVQSLALF TPALFPRSLD LSTDPVLATE KINRFIKAVT GWKMNLPLEG VSTDSTLLFN TYVHFQGTMR GFSQLPGVHE FWVDNSISVS VPMISGTGNF QHWSDAQNNF SVTCVPLGER ATLLLIQPHC TSDLDRVEAL IFRNDLLTWI ENPPPRAIRL TLPQLEIRGS YNLQDLLAED KLPTLLGAEA NLSNIGDTNP RVGEVLNSIL LELKAGEEEQ PTTSVQQPGS PEALDVTLSS PFLFAIYEQD SGTLHFLGRV NNPQSVV

### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against AGT. It has been selected for its ability to recognize AGT in immunohistochemical staining and western blotting.


### [APPLICATIONS]


Designed by Cloud-Clone Corp., Assembled by Uscn Life Science Inc. ISO9001:2008; ISO13485:2003; C € 11271 Richmond Avenue Suite H104, Houston, TX 77082, USA | Toil free: 001-888-960-7402 | Fax: 001-832-538-0088 | Http://www.cloud-clone.us | E-mail: mail@cloud-clone.us Export Processing Zone Building F, Wuhan, Hubel 430056, PRC | Toil free: 0086-800-880-0687 | Fax: 0086-27-8425-8551 | Http://www.uscnk.com | E-mail: mail@uscnk.com

# Coud-Clone Corp.

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

### [ <u>IMAGES</u> ]





Used in DAB staining on fromalin fixed paraffin- embedded Liver tissue

# paraffin- embedded Liver tis

Used in Western Blot ,Sample: Recombinant AGT,Mouse

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant AGT (Ala23~Val477) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

### [ <u>STORAGE</u> ]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.