# Coud-Clone Corp.

PAA816Mu01 Polyclonal Antibody to Alpha-1-Acid Glycoprotein (a1AGP) Organism Species: Mus musculus (Mouse) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [PRODUCT INFORMATION]

Immunogen: a1AGP, Mouse Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant a1AGP (Gln19~Ala207) expressed in E.coli.

USCN Accession No.: RPA816Mu01

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-QN PEHANFTIGE PITNETLSWL SDKWFFMGAA FRKLEYRQAI QTMQSEFFYL TTNLINDTIE LRESQTIGDQ CVYNSTHLGF QRENGTFSKY EGGVETFAHL IVLRKHGAFM LAFDLKDEKK RGLSLYAKRP DITPELREVF QKAVTHVGMD ESEIIFVDWK KDRCGQQEKK QLELGKETKK DPEEGQA

#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against a1AGP. It has been selected for its ability to recognize a1AGP in immunohistochemical staining and western blotting.

## [APPLICATIONS]


Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100

# Coud-Clone Corp.

Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

## [<u>IMAGES</u>]



Western Blot kDa 70 44 33 26 22 ←28kDa ←24kDa 18 14

Used in DAB staining on fromalin fixed paraffin- embedded Liver tissue [ CONTENTS ] Used in Western Blot ,Sample: Recombinant a1AGP,Mouse

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [QUALITY CONTROL]

**Content:** The quality control contains recombinant a1AGP (GIn19~Ala207) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

## [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.