PAA835Ra01 Polyclonal Antibody to Corticotropin Releasing Hormone (CRH) Organism Species: Rattus norvegicus (Rat) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Coud-Clone Corp.

11th Edition (Revised in May, 2016)

[PROPERTIES]

Source: Polyclonal antibody preparation Host: Rabbit Purification: Antigen-specific Affinity Chromatography. Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

[IMMUNOGEN]

Immunogen: RPA835Ra01-Recombinant Corticotropin Releasing Hormone (CRH)

[APPLICATIONS]

Western blotting: 0.5-2ug/ml Immunocytochemistry in formalin fixed cells: 5-20ug/ml Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. Western Blot A. Sample: Recombinant CRH, Rat




Figure 2. DAB staining on IHC-P Samples: A. Rat Testis Tissue

B. Rat Spleen Tissue

1304 Langham Creek Dr. Soire 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.us | mail@cloud-close.as Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0687 | www.cloud-close.com | mail@cloud-close.com