

PAB837Ra02

Polyclonal Antibody to Interleukin 2 Receptor Alpha (IL2Ra)
Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PRODUCT INFORMATION ]

Immunogen: IL2Ra, Rat

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM:** 100µg



Sample: Recombinant IL2Ra, Rat

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant IL2Ra (Leu23~Gln235) expressed in E.coli.

Accession No.: RPB837Ra02

 $\textbf{Sequence:} \ \ \textbf{The target protein is fused with two N-terminal Tags, His-tag and}$ 

S-tag and its sequence is listed below.

MHHHHHHSSG LVPRGSGMK ETAAAKFERQH MDSPDLGTDDD DKAMADIGSEF-LCLYDPPE VPNATFKALS YKNGTILNCE CKRGFRRLNE LVYMACLGNS WSNNCQCTSN SHDNSREQVT PQPEGQKEQQ TTDTQKSTQS VYQENLAGHC REPPPWRHED TKRIYHFVEG QIVLYTCIQG YKALQRGPAI SICKTVCGEI RWTHPQLTCV DEKEHHQFLA SFESQGSRNS FPESFASCPT PNTDESQLTE ATTTMETEVE TKEYQ



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against IL2Ra. It has been selected for its ability to recognize IL2Ra in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant IL2Ra (Leu23~Gln235) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.