

PAK204Hu01

Polyclonal Antibody

To A Disintegrin And Metalloproteinase With Thrombospondin 4 (ADAMTS4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: ADAMTS4, Human **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

Immunoglobulin Type: IgG **UOM**: 100μg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant ADAMTS4 (Thr437~Pro575) expressed in *E.coli*.

USCN Accession No.: RPK204Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-TFPG KDYDADRQCQ LTFGPDSRHC PQLPPPCAAL WCSGHLNGHA MCQTKHSPWA DGTPCGPAQA CMGGRCLHMD QLQDFNIPQA GGWGPWGPWG DCSRTCGGGV OFSSRDCTRP VPRNGGKYCE GRRTRFRSCN TEDCP

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against ADAMTS4. It has been selected for its ability to recognize ADAMTS4 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200




Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on fromalin fixed paraffin- embedded Lung tissue



Used in Western Blot ,Sample: Recombinant ADAMTS4,Human

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant ADAMTS4 (Thr437~Pro575) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.