RPB975Ra01 50µg Recombinant Tenascin C (TNC) Organism Species: Rattus norvegicus (Rat) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Coud-Clone Corp.

#### [PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Asp513~Ser621

Tags: N-terminal His-Tag

Tissue Specificity: Heart.

**Purity: >92%** 

Traits: Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

**Applications:** SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 12.6kDa

Accurate Molecular Mass: 18kDa as determined by SDS-PAGE reducing conditions.

#### Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

## [ <u>USAGE</u> ]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.



#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [SEQUENCE]

DGQCICED GFTGSDCAEL SCPGDCHGHG RCVNGQCICH EGFTGXDCKE QRCPSDCHGQ GRCEDGQCIC HEGFTGLDCG QRSCPNDCSN QGQCVAGRCI CNEGYTGVDC S

#### [IDENTIFICATION]




Figure 1. Gene Sequencing (Extract)

# Coud-Clone Corp.

	kDa 70
. 3	44
	33
	26
	22
-	18
-	14
3.4	10

Figure 2. SDS-PAGE