

TAX656Ge22
Anti-Hexa Histidine (H6) Tag Monoclonal Antibody
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



#### [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG

**Purification:** Protein A/G Affinity Chromatography.

Clone number: D4

Traits: Liquid

Concentration: 500µg/mL

**UOM**: 200µg

**Applications:** WB; ICC/IF; Dot Blot; ELISA; IP; EMSA; Purification.

# [ IMMUNOGEN ]

**Immunogen:** Synthetic Peptide, H6 conjugated to OVA.

Accession No.: CPX656Ge21

## [ APPLICATIONS ]

Western blotting: 0.5-5ug/ml

Immunocytochemistry in formalin fixed cells/Immunofluorescence: 5-30ug/ml

Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were



observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. Western Blot
Sample: Recombinant Protein RPB693Mu01 with
His-tag