


Phospho-Vimentin (Ser83) Ab

| Cat.#: AF2433 Size: 100ul,200ul | Concn.: 1mg/ml Source: Rabbit | Mol.Wt.: 57kDa Clonality: Polyclonal |
|------------------------------------|---|---|
| Application: | WB 1:500-1:2000, IHC 1:50-1:200 | |
| Reactivity: | Human,Mouse,Rat | |
| Purification: | The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns. | |
| Specificity: | Phospho-Vimentin (Ser83) Ab detects endogenous levels of Vimentin. | |
| Immunogen: | A synthesized peptide derived from human Vimentin around the phosphorylation site of Ser83. | |
| Uniprot: | P08670 | |
| Subcellular Location: | Cytoplasm. | |
| Tissue Specificity: | Highly expressed in fibroblasts, some expression in T- and B- lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone- independent mammary carcinoma cell lines. | |
| Similarity: | The central alpha-helical coiled-coil IF rod domain mediates elementary homodimerization.The [IL]-x-C-x-x-[DE] motif is a proposed target motif for cysteine S-nitrosylation mediated by the iNOS-S100A8/A9 transnitrosylase complex.Belongs to the intermediate filament family. | |
| Storage Condition and Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt | |



Western blot analysis of extracts of Hela cells treated with Paclitaxel, using Phospho-Vimentin (Ser83) Ab.





AF2433 at 1/100 staining Mouse lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.