

Phospho-Catenin-β (Ser37) Ab

Cat.#: AF3266 Size: 100ul,200ul	Concn.: 1mg/ml Source: Rabbit	Mol.Wt.: 92kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500	
Reactivity:	Human,Mouse,Rat	
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.	
Specificity:	Phospho-Catenin- β (Ser37) Ab detects endogenous levels of Catenin- β only when phosphorylated at Serine 37	
Immunogen:	A synthesized peptide derived from human Catenin- β around the phosphorylation site of Serine 37	
Uniprot:	P35222	
Description:	Beta-catenin is an adherens jur junctions (AJs; also called the ze for the establishment and main such as those lining organ surfa between cells, communicate a are present, and anchor the act these roles, AJs regulate norma	onula adherens) are critical itenance of epithelial layers, aces. AJs mediate adhesion signal that neighboring cells tin cytoskeleton. In serving
Subcellular Location:	Cytoplasm. Nucleus. Cytoplasm > adherens junction. Cell juncti Cytoplasmic when it is unstabil phosphorylation) or bound to C nucleus when it is stabilized (lo Interaction with GLIS2 and MUC translocation. Interaction with B localization.	on. Cell membrane. ized (high level of DH1. Translocates to the w level of phosphorylation). 1 promotes nuclear
Tissue Specificity:	Expressed in several hair follicl peripheral matrix cells, and cel sheaths. Expressed in colon. Pr protein level). Expressed in bre level) (PubMed:29367600).	ls of the outer and inner root esent in cortical neurons (at
Similarity:	Belongs to the beta-catenin family.	
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt	



kDa 1 2


250-

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

Western blot analysis of Catenin- β phosphorylation expression in 293 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.

150 -70 -35 - - - - + Peotide

Western blot analysis of extracts of various tissue sample, using Phospho-Catenin- β (Ser37) Ab.



AF3266 at 1/200 staining human colon cancer tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat antirabbit Ab was used as the secondary.



AF3266 staining SW626 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.