Phospho-Catenin beta (Thr41/Ser45) Ab

Cat.#: AF3268 Concn.: 1mg/ml Mol.Wt.: 92kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-Catenin- beta (Thr41/Ser45) Ab detects

endogenous levels of Catenin- beta only when phosphorylated at Threonine 41/Serine 45

Immunogen: A synthesized peptide derived from human Catenin- beta

around the phosphorylation site of Threonine 41/Serine 45

Uniprot: P35222

Description: Beta-catenin is an adherens junction protein. Adherens

junctions (AJs; also called the zonula adherens) are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces. AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior.

Subcellular Location: Cytoplasm. Nucleus. Cytoplasm > cytoskeleton. Cell junction

> adherens junction. Cell junction. Cell membrane.
Cytoplasmic when it is unstabilized (high level of phosphorylation) or bound to CDH1. Translocates to the nucleus when it is stabilized (low level of phosphorylation).
Interaction with GLIS2 and MUC1 promotes nuclear translocation. Interaction with EMD inhibits nuclear

localization.

Tissue Specificity: Expressed in several hair follicle cell types: basal and

peripheral matrix cells, and cells of the outer and inner root sheaths. Expressed in colon. Present in cortical neurons (at protein level). Expressed in breast cancer tissues (at protein

level) (PubMed:29367600).

Similarity: Belongs to the beta-catenin family.

Storage Condition and

Buffer:


Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20

°C.Stable for 12 months from date of receipt



Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis of Catenin- beta phosphorylation expression in SW626 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.

Western blot analysis of Phospho-Catenin beta (Thr41/Ser45) expression in various lysates



AF3268 staining SW626 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.