


Phospho-14-3-3 zeta/ delta (Thr232) Ab

Cat.#: AF3357 Size: 100ul,200ul	Concn.: 1mg/ml Source: Rabbit	Mol.Wt.: 28kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500	
Reactivity:	Human,Mouse,Rat	
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.	
Specificity:	Phospho-14-3-3 zeta/ delta (Thr232) Ab detects endogenous levels of 14-3-3 zeta/ delta only when phosphorylated at Threonine 232	
Immunogen:	A synthesized peptide derived f delta around the phosphorylatic	
Uniprot:	P63104	
Description:	14-3-3 zeta is a protein of the 1 which mediate signal transducti phosphoserine-containing prote	ion by binding to
Subcellular Location:	Cytoplasm. Melanosome. Locate melanosomes.	ed to stage I to stage IV
Similarity:	Belongs to the 14-3-3 family.	
Storage Condition and Buffer:	Rabbit lgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt	



Western blot analysis of 14-3-3 zeta/ delta phosphorylation expression in UV treated Jurkat whole cell lysates,The lane on the left is treated with the antigen-specific peptide.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com




Western blot analysis of Phospho-14-3-3 zeta/ delta (Thr232) Ab expression in UV treated Jurkat cells lysates. The lane on the right is treated with the antigen-specific peptide.



AF3357 at 1/100 staining human brain tissues sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at $39^{\circ}C$



AF3357 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.



AF3357 at 1/100 staining Rat brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.