


Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

Phospho-ANXA2 (Tyr24) Ab

Cat.#: AF7096 Size: 50ul,100ul,200ul	Concn.: 1mg/ml Source: Rabbit	Mol.Wt.: 37kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200	
Reactivity:	Human,Mouse,Rat	
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.	
Specificity:	Phospho-ANXA2 (Tyr24) Ab detects endogenous levels of ANXA2 only when phosphorylated at Tyr24	
Immunogen:	A synthesized peptide derived from human ANXA2 around the phosphorylation site of Tyr24	
Uniprot:	P07355	
Subcellular Location:	Secreted > extracellular space > extracellular matrix > basement membrane. Melanosome. In the lamina beneath the plasma membrane. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Translocated from the cytoplasm to the cell surface through a Golgi-independent mechanism.	
Similarity:	A pair of annexin repeats may form one binding site for calcium and phospholipid.Belongs to the annexin family.	
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt	



Western blot analysis of Phospho-ANXA2 (Tyr24) in lysates of HepG2, using Phospho-ANXA2 (Tyr24) Ab(AF7096).





AF7096 at 1/100 staining human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF7096 at 1/100 staining rat heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.