Phospho-PSD95 (Tyr609) Ab

Cat.#: AF7338 Mol.Wt.: 105kDa Concn.: 1mg/ml Size: 50ul.100ul.200ul Source: Rabbit Clonality: Polyclonal

WB 1:500-1:2000 Application:

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-PSD95 (Tyr609) Ab detects endogenous levels of

PSD95 only when phosphorylated at Tyr609

A synthesized peptide derived from human PSD95 around Immunogen:

the phosphorylation site of Tyr609

P78352 Uniprot:

Subcellular Location: Cell membrane. Cell junction, synapse, postsynaptic cell

> membrane, postsynaptic density. Cell projection, axon. Cell junction, synapse. High levels in postsynaptic density of neurons in the forebrain. Also in presynaptic region of inhibitory synapses formed by cerebellar basket cells on

axon hillocks of Purkinje cells.

Brain. Tissue Specificity:

Similarity: The PDZ domain 3 mediates interaction with ADR1B. The L27

domain near the N-terminus of isoform 2 is required for

HGS/HRS-dependent targeting to postsynaptic


density. Belongs to the MAGUK family.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt



Western blot analysis of extracts of human brain tissue sample, using Phospho-PSD95 (Tyr609) Ab(AF7338).



w/v milk , 1X TBS, 0.1% Tween@20 at $4^{\circ}C$ with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.