

ATAGENIX LABORATORIES

Catalog Number:ATMA10324Mo Anti SARS-CoV-2(2019-nCoV) Nucleocapsid antibody(54#)

Product Details

Summary

Product name Anti SARS-CoV-2(2019-nCoV) Nucleocapsid antibody(54#)

descriptionThis antibody was produced from a hybridoma resulting from the fusion of a

mouse myeloma with B cells obtained from a mouse immunized with purified,

recombinant SARS-CoV-2(2019-nCoV) Nucleocapsid protein.

Accession # P0DTC9

Alternative names Nucleoprotein, N, NC, Protein N

Purity Nucleoprotein, N, NC, Protein N

Stability &StorageUse a manual defrost freezer and avoid repeated freeze thaw cycles.

Store at 2 to 8 °C for one week .

Store at -20 to -80 °C for twelve months from the date of receipt.

Spcificty Recognizes SARS-CoV-2 Nucleocapsid protein

lsotype lgG 1

Host Mouse

Clonality Monoclonal

Clone No. 54#

Conjugation Unconjugated

Species reactivity Severe acute respiratory syndrome coronavirus 2 (2019-nCoV) (SARS-CoV-2)

Tested applications ELISA

Immunogen Recombinant SARS-CoV-2 Nucleocapsid protein(Met1-Ala419)

Background

Coronavirus N protein is required for coronavirus RNA synthesis, and has RNA chaperone activity that may be involved in template switch. Nucleocapsid protein is a most abundant protein of coronavirus. N protein packages the positive strand viral genome RNA into a helical ribonucleocapsid (RNP) and plays a fundamental role during virion assembly through its interactions with the viral genome and membrane protein M. Plays an important role in enhancing the efficiency of subgenomic viral RNA transcription as well as viral replication. Because of the conservation of N protein sequence and its strong immunogenicity, the N protein of coronavirus is chosen as a diagnostic tool.

Product performance



ATAGENIX LABORATORIES

Catalog Number:ATMA10324Mo Anti SARS-CoV-2(2019-nCoV) Nucleocapsid antibody(54#)

Form Liquid

Buffer PBS, pH7.4, containing 0.05% proclin300, 50% glycerol.

Concentration 1.18mg/ml

MW 45kDa

Application

Dilution Range Elisa:1:200-1:2000

Note

For research use only .Not for use in clinical diagnostic procedures.