

MMP3 Ab

Cat.#: BF0526
Size: 50ul,100ul,200ul

Concn.: 1mg/ml
Source: Mouse

Mol.Wt.: 54kDa
Clonality: Monoclonal

Application:	ELISA 1/10000, WB 1/500 - 1/2000, FCM 1/200 - 1/400
Reactivity:	Human
Purification:	Affinity-chromatography.
Specificity:	MMP3 Ab detects endogenous levels of total MMP3.
Immunogen:	Purified recombinant fragment of human MMP3 expressed in E. Coli.
Uniprot:	P08254
Description:	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.
Subcellular Location:	Secreted > extracellular space > extracellular matrix.
Similarity:	The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.Belongs to the peptidase M10A family.
Storage Condition and Buffer:	Mouse IgG1 in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.





Figure 1: Western blot analysis using MMP3 mAb against human MMP3 (AA: 189-441) recombinant protein. (Expected MW is 54 kDa)



Flow cytometric analysis of NIH/3T3 cells using MMP3 mouse mAb (green) and negative control (red).

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.