ABCD2 Antibody

Cat.#: DF9250 Concn.: 1mg/ml Mol.Wt.: 83 kDa Size: 100ul.200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:1000-3000, IF/ICC 1:100-1:500, ELISA(peptide)

1:20000-1:40000

*The optimal dilutions should be determined by the end

user.

Reactivity: Human, Mouse

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: ABCD2 Antibody detects endogenous levels of total ABCD2.

A synthesized peptide derived from human ABCD2, Immunogen:

corresponding to a region within the internal amino acids.

Uniprot: Q9UBJ2

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from Mouse testis, using ABCD2 Antibody. The lane on the left was treated with


blocking peptide.

Western blot analysis of ABCD2 using HeLa whole cell lysates



Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com



DF9250 staining 3T3 by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF9250 1:200) and mouse antibeta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at 4° C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.