ABCF3 Antibody

Cat.#: DF9251 Concn.: 1mg/ml Mol.Wt.: 80 kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:1000-3000, IF/ICC 1:100-1:500, ELISA(peptide)

1:20000-1:40000

*The optimal dilutions should be determined by the end

user.

Reactivity: Human, Mouse, Rat

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: ABCF3 Antibody detects endogenous levels of total ABCF3.

Immunogen: A synthesized peptide derived from human ABCF3,

corresponding to a region within the internal amino acids.


Uniprot: Q9NUQ8

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis of ABCF3 using K562 whole cell lysates

DF9251 staining HepG2 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF9251 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.