

## BNIPL Antibody

Cat.#: DF9267  
Size: 100ul,200ul

Concn.: 1mg/ml  
Source: Rabbit

Mol.Wt.: 40 kDa  
Clonality: Polyclonal

**Application:** WB 1:1000-3000, IHC 1:200, ELISA(peptide)  
1:20000-1:40000  
\*The optimal dilutions should be determined by the end user.

**Reactivity:** Human,Monkey


**Purification:** The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

**Specificity:** BNIPL Antibody detects endogenous levels of total BNIPL.


**Immunogen:** A synthesized peptide derived from human BNIPL, corresponding to a region within the internal amino acids.

**Uniprot:** Q7Z465

**Storage Condition and Buffer:** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of BNIPL using COS7 whole cell lysates



DF9267 at 1/100 staining human lung cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.