

MAA519Hu26 Monoclonal Antibody to Apolipoprotein A1 (APOA1) Organism Species: Homo sapiens (Human) *Instruction manual* 

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Suite 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.us | mail@cloud-clone.as Export Processing Zone, Wuhan, Hubei 436056, PRC | 0086-000-880-0687 | www.cloud-clone.com | mail@cloud-clone.com

# Cloud-Clone Corp.

# [PROPERTIES]

Source: Monoclonal antibody preparation Host: Mouse Antibody isotype: IgG2b Kappa Purification: Protein A/G Affinity Chromatography. Clone number: C10 Traits: Liquid Concentration: 500µg/mL UOM: 200µg Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

#### [IMMUNOGEN]

Immunogen: Recombinant APOA1 (Gln122~Gln267) expressed in *E.coli*. Accession No.: RPA519Hu01

# [APPLICATIONS]

Western blotting: 0.5-5ug/ml Immunocytochemistry in formalin fixed cells: 5-30ug/ml Immunohistochemistry in formalin fixed frozen section: 5-30ug/ml Immunohistochemistry in paraffin section: 5-30ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

# [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

# [QUALITY CONTROL]

**Content:** The quality control contains recombinant APOA1 disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine (DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.

#### [STORAGE AND STABILITY]

Coud-Clone Corp.


Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [IDENTIFICATION]



#### Figure 1. Western Blot

A. Sample: Recombinant APOA1, Human

B. Human Serum

Primary Ab: 2µg/mL Mouse Anti-Human APOA1 Ab Second Ab: 1:5000 Dilution of HRP-Linked Rabbit Anti-Mouse IgG Ab (Catalog: SAA544Mu09)

1304 Langham Creek Dr, Suite 226, Huustun, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.us | mail@cloud-cloue.us Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0687 | www.cloud-clone.com | mail@cloud-clone.com

# Coud-Clone Corp.




Figure 2. DAB staining on IHC-P Samples:

- A. Human Liver Tissue
- B. Human Kidney Tissue

1304 Langham Creek Dr. Saire 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.ox | mail@cloud-close.ax Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-008-0687 | www.cloud-close.com | mail@cloud-close.com