

#### **PAA011Mi01**

Polyclonal Antibody to Brain Derived Neurotrophic Factor (BDNF)

**Organism Species: Homo sapiens (Human)** 

Mus musculus (Mouse)

Rattus norvegicus (Rat)

**Oryctolagus cuniculus (Rabbit)** 

Sus scrofa; Porcine (Pig)

Capra hircus; Caprine (Goat)

Equus caballus; Equine (Horse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



#### [PROPERTIES]

Source: Polyclonal antibody preparation

**Host:** Rabbit

**Purification:** Antigen-specific Affinity Chromatography.

Traits: Liquid

Concentration: 200µg/mL

**UOM**: 100µg

Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

# [ IMMUNOGEN ]

Immunogen: Recombinant BDNF (Pro20~Arg252) expressed in *E.coli*.

Accession No.: RPA011Mi01

# [ORGANISM SPECIES MORE]

**React with:** Human, mouse, rat, rabbit, porcine, caprine, equine;

Other species have not been detected.

### [APPLICATIONS]

Western blotting: 0.5-2ug/ml

Immunocytochemistry in formalin fixed cells: 5-20ug/ml

Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml

Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

# [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant BDNF disposed in loading buffer.



Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

Loading Buffer: 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol,

0.02% BPB, 50mM DTT and 0.02% NaN $_{\!3}.$ 

# [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [IDENTIFICATION]




Figure 1. Western Blot Sample: Recombinant BDNF

# Coud-Clone Corp.




Figure 2. DAB staining on IHC-P

Samples:

**Mouse Ovary Tissue**