

PAA303Ra01

Polyclonal Antibody to Galectin 3 (GAL3)
Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: GAL3, Rat **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

Immunoglobulin Type: IgG **UOM**: 100μg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant GAL3 (Met1~Ile262) expressed in *E.coli*.

Accession No.: RPA303Ra01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSG S-MADGFSLNDA LAGSGNPNPQ GWPGAWGNQP GAGGYPGASY PGAYPGQAPP GGYPGQAPPS AYPGPTGPSA YPGPTAPGAY PGPTAPGAFP GQPGGPGAYP SAPGAYPSAP GAYPATGPFG APTGPLTVPY DMPLPGGVMP RMLITIIGTV KPNANSITLN FKKGNDIAFH FNPRFNENNR RVIVCNTKQD NNWGREERQS AFPFESGKPF KIQVLVEADH FKVAVNDVHL LQYNHRMKNL REISQLGIIG DITLTSASHA MI

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against GAL3. It has been selected for its ability to recognize GAL3 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500




Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on fromalin fixed paraffin- embedded brain tissue



Used in Western Blot ,Sample:

Recombinant GAL3.Rat

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant GAL3 (Met1~Ile262) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.