

PAA656Hu01

Polyclonal Antibody to Diamine Oxidase (DAO)

**Organism Species: Homo sapiens (Human)** 

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

## [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific Affinity Chromatography.

Traits: Liquid

Concentration: 200µg/mL

**UOM**: 100µg

Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

### [ IMMUNOGEN ]

Immunogen: RPA656Hu01-Recombinant Diamine Oxidase (DAO)

## [APPLICATIONS]

Western blotting: 0.5-2ug/ml

Immunocytochemistry in formalin fixed cells: 5-20ug/ml

Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml

Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.



## [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]




Figure 1. Western Blot

A. Sample: Recombinant DAO, Human

**B. Porcine Kidney Tissue** 

Primary Ab: 3µg/mL Rabbit Anti-Human DAO Ab

Second Ab: 1:2000 Dilution of HRP-Linked Guinea pig

Anti-Rabbit Ab (Catalog: SAA544Rb59)




Figure 2. DAB staining on

IHC-P

Samples:

A. Human Kidney Tissue