PAJ226Hu02 Polyclonal Antibody to Biglycan (BGN) Organism Species: Homo sapiens (Human) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

#### [PRODUCT INFORMATION]

Cloud-Clone Corp.

Immunogen: BGN, Human Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

#### [ IMMUNOGEN INFORMATION ]

**Immunogen:** Recombinant BGN (Thr230~Tyr366) with N-terminal His-Tag expressed in *E.coli*.

Accession No.: RPJ226Hu02

#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against BGN. It has been selected for its ability to recognize BGN in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-5000 Optimal working dilutions must be determined by end user.



## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [QUALITY CONTROL]

**Content:** The quality control contains recombinant BGN (Thr230~Tyr366) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.


5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.


# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

## [<u>IMAGES</u>]



Used in Western Blot, Sample: Recombinant BGN, Human




Used in DAB staining on fromalin fixed paraffin- embedded liver tissue

1304 Langham Creek Dr, Suite 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-clone.us | mail@cloud-clone.us Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-800-880-0687 | www.cloud-clone.com | mail@cloud-clone.com C <u>Cloud-Clone Corp</u>.

PAJ226Hu02 Polyclonal Antibody to Biglycan (BGN) Organism Species: Homo sapiens (Human) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



Used in Western Blot, Sample: Lane1: Porcine Spleen Tissue Lane2: Porcine Kidney Tissue Primary Ab: 5µg/mL Rabbit Anti-Human BGN Ab Second Ab: 1:1000 Dilution of HRP-Linked Guinea pig Anti-Rabbit Ab (Catalog: SECab05)