

RPA133Po01 10µg

**Recombinant Tumor Necrosis Factor Alpha (TNFa)** 

Organism Species: Sus scrofa; Porcine (Pig)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



#### [PROPERTIES]

**Source:** Prokaryotic expression.

Host: E. coli

Residues: Leu77~Leu232 Tags: N-terminal His-Tag Tissue Specificity: Liver.

Subcellular Location: Cell membrane; Secreted Single-pass type II membrane

protein.

**Purity:** >98%

**Traits:** Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive

Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.8

Predicted Molecular Mass: 20.7kDa

Accurate Molecular Mass: 21kDa as determined by SDS-PAGE reducing conditions.

## [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [SEQUENCE]

LRSS SQTSDKPVAH VVANVKAEGQ LQWQSGYANA LLANGVKLKD NQLVVPTDGL YLIYSQVLFR GQGCPSTNVF LTHTISRIAV SYQTKVNLLS AIKSPCQRET PEGAEAKPWY EPIYLGGVFQ LEKDDRLSAE INLPDYLDFA ESGQVYFGII AL

# [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE