

RPF182Hu01 50µg
Recombinant 17-Beta-Hydroxysteroid Dehydrogenase Type 12 (HSD17b12)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Met1~Asn312

Tags: N-terminal His-Tag

Tissue Specificity: Ovary, Breast.

Subcellular Location: Endoplasmic reticulum membrane; Multi-pass membrane protein.

Purity: >98%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.3

Predicted Molecular Mass: 38.0kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MESALPAAGF LYWVGAGTVA YLALRISYSL FTALRVWVG NEAGVGPGLG
EWAVVTGSTD GIGKSYAEEL AKHGMKVLI SRSKDKLDQV SSEIKEKFKV
ETRTIAVDFA SEDIYDKIKT GLAGLEIGIL VNNVGMSY EY PEYFLDVPDL
DNVIKKMINI NILSVCKMTQ LVLPGMVERS KGAILNISSG SGMLPVPLLT
IYSATKTFVD FFSQCLHEEY RSKGVFVQSV LPYFVATKLA KIRKPTLDKP
SPETFVKS AI KTVGLQSR TN GYLIHALMGS IISNLPSWIY LKIVMNMNKS
TRAHYLKTK KN

[IDENTIFICATION]




Figure 1. SDS-PAGE