ో <u>Cloud-Clone Corp</u>.

RPH574Hu01 100µg Recombinant N-Acetylgalactosaminidase Alpha (NAGa) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Leu18~Asp217 (Accession # P17050), with N-terminal His-Tag. Host: *E. coli* Subcellular Location: Lysosome. Purity: >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. Predicted isoelectric point: 5.1 Predicted Molecular Mass: 24.1kDa Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[<u>USAGE</u>]

Reconstitute in ddH₂O.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF- LDN GLLQTPPMGW LAWERFRCNI NCDEDPKNCI SEQLFMEMAD RMAQDGWRDM GYTYLNIDDC WIGGRDASGR LMPDPKRFPH GIPFLADYVH SLGLKLGIYA DMGNFTCMGY PGTTLDKVVQ DAQTFAEWKV DMLKLDGCFS TPEERAQGYP KMAAALNATG RPIAFSCSWP AYEGGLPPRV NYSLLADICN LWRNYDD

[REFERENCES]

- 1. Wang A.M., et al. (1990) J. Biol. Chem. 265:21859-21866.
- 2. Wang A.M., Desnick R.J. (1991) Genomics 10:133-142.
- 3. Tsuji S., et al. (1989) Biochem. Biophys. Res. Commun. 163:1498-1504.
- 4. Yamauchi T., et al. (1990) Biochem. Biophys. Res. Commun. 170:231-237.