

RPH918Hu01 50µg

Recombinant Hephaestin (HEPH)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Ala24~Cys366 Tags: N-terminal His-Tag

Tissue Specificity: Breast, Colon.

Subcellular Location: Membrane, Single-pass type I membrane protein.

Purity: >95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive

Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.4

Predicted Molecular Mass: 42.1kDa

Accurate Molecular Mass: 42kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

ATRVYYL GIRDVQWNYA PKGRNVITNQ
PLDSDIVASS FLKSDKNRIG GTYKKTIYKE YKDDSYTDEV AQPAWLGFLG
PVLQAEVGDV ILIHLKNFAT RPYTIHPHGV FYEKDSEGSL YPDGSSGPLK
ADDSVPPGGS HIYNWTIPEG HAPTDADPAC LTWIYHSHVD APRDIATGLI
GPLITCKRGA LDGNSPPQRQ DVDHDFFLLF SVVDENLSWH LNENIATYCS
DPASVDKEDE TFQESNRMHA INGFVFGNLP ELNMCAQKRV AWHLFGMGNE
IDVHTAFFHG QMLTTRGHHT DVANIFPATF VTAEMVPWEP GTWLISCQVN
SHFRDGMQAL YKVKSC

[IDENTIFICATION]




Figure 1. SDS-PAGE